

Optimal Iodine Nutrition during Pregnancy, Lactation and the Neonatal Period

Delange F.

International Council for Control of Iodine Deficiency Disorders (ICCIDD); Department of Pediatrics, University of Brussels, Brussels, Belgium

Iodine of maternal origin is required for brain development of the progeny during fetal and early postnatal life. Therefore, the iodine requirements of the mother are increased during pregnancy and lactation. This paper reevaluates the iodine requirements during pregnancy, lactation and the neonatal period and formulates original proposals for the median concentrations of urinary iodine indicating optimal iodine nutrition during these three critical periods of life. Based on an extensive and critical review of the literature on thyroid physiopathology during the perinatal period, the following proposals are made: the iodine requirements are 250-300 µg/day during pregnancy, 225-350 µg/day during lactation and 90 µg/day during the neonatal period. The median urinary iodine indicating optimal iodine nutrition during these three periods should be in the range 150-230 µg/L. These figures are higher than those recommended so far by international agencies.

Key Words: Iodine, Nutrition, Pregnancy, Lactation, Neonatal Period, Median urinary iodine

Introduction

The thyroid economy undergoes a series of metabolic changes during pregnancy and lactation.¹⁻⁴ One of the factors involved in these

changes is the increased requirement of iodine in the mother due to the transfer of thyroxine (T₄) and of iodide from mother to fetus during pregnancy and to the loss of iodide in breast milk during lactation. These two processes are required in order to ensure normal brain development and prevention of mental retardation in the offspring.⁵⁻¹⁰

The objectives of this paper are:

1. To review the data from the literature on the iodine requirements during pregnancy, lactation and the neonatal period.
2. To offer practical recommendations regarding the median concentrations of urinary iodine indicating optimal iodine nutrition during these critical periods of life.

Requirement of Iodine During Pregnancy and Lactation

The requirement of iodine is increased during pregnancy because of at least three factors: 1) There is an increased requirement of T₄ in order to maintain a normal global metabolism in the mother. 2) There is a transfer of T₄ and iodide from the mother to the fetus and 3) There is supposed to be an increased loss of iodide through the kidney due to an increase in the renal clearance of iodide.

Correspondence: Professor François Delange, MD, PhD, 153, avenue de la Fauconnerie B-1170, Brussels, Belgium

E-mail: fdelange@ulb.ac.be

Because of these three factors, the recommended dietary intake of iodine during pregnancy is higher than the value of 150 µg/day recommended for non-pregnant adults and adolescents.^{11,12} Below this critical threshold of 150 µg/day, the iodine balance is negative during pregnancy.¹³ WHO/UNICEF/ICCIDD recommend an iodine intake of 200 µg/day for pregnant women,¹¹ i.e. a percentage increase of 33% over non-pregnant women. The Institute of Medicine (IOM) of the US Academy of Sciences recommends a higher intake of 220 µg/day,¹² i.e. an increase of some 47%, and other organizations recommend 175 to 230 µg/day.^{14,15}

Increase in the T₄ requirements

The daily requirement of T₄ in order to maintain euthyroidism in hypothyroid women increases by 10 to 150% during pregnancy with a median increment of 40-50%.¹⁶⁻¹⁸ This represents an additional dose of 75 to 150 µg T₄/day, i.e. 50 to 100 µg iodine.

Transfer of T₄ and iodide from mother to fetus

The transfer of T₄ from mother to fetus, including before the onset of fetal thyroid function, is not quantified but it has been estimated that up to 40% of the T₄ measured on cord at birth is still of maternal origin.⁸

The transfer of iodide is also difficult to quantify but considering that the iodine content of the fetal thyroid increases progressively from less than 2 µg at 17 weeks of gestation¹⁹ up to 300 µg at term,²⁰⁻²³ that the T₄ iodine at term probably averages 500 µg²⁴ and that the substitutive dose of T₄ in hypothyroid neonates is 50-75 µg/day,^{25,26} it can be estimated that the transfer of iodide from mother to fetus represents some 50 µg/day. It has been estimated at 75 µg/day by the IOM.¹²

Increased renal clearance of iodide

It is often stated that the increase in iodine requirement during pregnancy is largely due to an increased loss of iodide through the

kidney because of an increased renal clearance of iodide. This should decrease the serum concentration of plasma inorganic iodide, PII.²⁷⁻³⁰ However, Liberman et al.³¹ showed on the contrary that there is no significant decline in the PII during pregnancy. In addition, as shown by the data collected in Table 1 and already by Dworkin et al.,¹³ almost all studies on urinary iodine during pregnancy showed that, in a given environment, the urinary excretion of iodide is almost similar in pregnant and non-pregnant women and in the general population, irrespective of the status of iodine nutrition in the population. Only the studies conducted by the group of Smyth et al.^{32,33} in Ireland, the United Kingdom and Sri Lanka, by Kung et al. in Hong Kong³⁴ and perhaps by Hess et al. in Switzerland³⁵ have shown a clear-cut increase in the urinary iodine excretion during pregnancy. The results reported for Switzerland by Brander et al.³⁶ are difficult to interpret because of the surprisingly low value of the urinary iodine in the general population reported in this study as Switzerland is known to be iodine sufficient.³⁷ On the contrary, some studies showed that urinary iodine decreases during gestation.³⁸⁻⁴⁰ It thus appears that the concept of systematically increased urinary loss of iodine during pregnancy is not firmly established.

Finally, it has to be underlined that no data are available on the possible storage and loss of iodide in the placenta itself.

Taking all these variables into consideration, it can be speculated that the additional requirement of iodine during pregnancy is at least 100-150 µg/day, i.e. an increment of almost 100% as compared to non-pregnant adults instead of the 33% proposed by WHO/UNICEF/ICCIDD.¹¹ Consequently, the requirement of iodine during pregnancy is at least 250 µg/day, probably in the range of 250 to 300 µg/day. This figure is still higher than the figure of 220 µg/day proposed by the IOM,¹² which did not take into account the increased requirement of T₄ during pregnancy.

Table 1. Comparison of the median or mean (in bold) urinary iodine ($\mu\text{g/L}$) in pregnant women and in the general population or in non-pregnant controls (publications 1990-2003)

Country	General population or controls	n	S/C	Pregnant women		Country	General population or controls	n	S/C	Pregnant women	
				Trimester	Urinary iodine					Trimester	Urinary iodine
Countries with no iodine deficiency						UK ³³	73 [†]	-	C	1	125 [†]
Chile ³¹	-	19	S	1	594*				C	2	170
				2	469			C	3	147	
				3	786						
				3 months PP	459	France ⁴⁰	50-80*	306	S	1	50 [†]
Iran ⁴³	193-312 [†]	403	C	1-3	186-338 [†]			224	S	3	54
Sweden ⁷⁷	-	51	S	1	180*	Belgium ³⁸	50-75*	334	C	1-2	50 [†]
				2	170					2-3	45
				3	145					1	56 [†]
Srilanka ³³	147 [†]	-	C	1	181 [†]			133	C	2	50
				2	136	Denmark ⁸³	50*	26	S	2	51 [†]
				3	154			26	S	3	40
USA ⁷⁸	130	290	C	1-3	148			26	S	1 week PP	30
Switzerland ³⁵ (2000)	115 [†]	511	C	2,3	138 [†]			26	S	26 weeks PP	50
Scotland ⁷⁹	138[†]	433	C	1	137[†]			26	S	52 weeks PP	58
Switzerland ³⁶ (1992)	91 [‡]	153	C	1-3	205*						
		31	C	1	267	Denmark ⁸⁴		-	C	5 days PP	40 [‡]
		56	C	2	206	Sudan ⁵⁵	76 [†]	47	S	3	38 [†]
		66	C	3	172			47	S	3 months PP	51
		15	S	1	325			47	S	6 months PP	30
		15	S	2	166			47	S	9 months PP	63
		15	S	3	183						
Iodine deficient countries						New Zealand ⁴⁹	24-47*	35	S	Monthly during pregnancy and 3, 6 and 12 months PP	24-52*
Singapore ^{34,80}	98 [†]	253	C	3	124 [†]						
		230	S	1	107 [†]						
		-		2	116						
		-		3	124						
		-		6 weeks PP	105						
		-		3 months PP	104	Italy ⁸⁵ (2002)	Marginal ID	67	C	1,2	74 [‡]
Sicily (Italy) ^{54,81}	46*	10	S	1,2,3	33*	Italy ⁸⁶ (1991)	Marginal ID	18	C	3	50*
Turkey ⁸²	85*	80	S	1-3	91*	Germany ⁸⁷	Mild ID	70	S	1	55 [‡]
Ireland ^{32,33}	70 [†]	38	S	1	135 [†]			70	S	11 days PP	50
				2	125						
				3	122	Hungary ⁸⁸	Mild ID	119	C	1,2,3	57 [‡]
		108	C	6 weeks PP	70						

n: number of subjects; S/C: Sequential (S) or cross-sectional study(C); 1,2,3: Trimesters of pregnancy; PP: Postpartum; ID: Iodine deficiency; 100 $\mu\text{g/L}$ = 0.78 $\mu\text{mol/L}$

* $\mu\text{g/day}$; [†] $\mu\text{g/L}$; [‡] $\mu\text{g/g creatinine}$

Table 2. Selective examples of the iodine content of breastmilk*

Countries	Medians or means ($\mu\text{g/L}$)
No iodine deficiency	
Korea	892
Japan	661
	33-385
USA	146
	168
	124
	145
	145
Sweden	93
	90
	70
Switzerland	78
Mild to moderate iodine deficiency	
Germany	93
	15-150
Belgium	95
France	82
	77
	74
	70
Spain	108
	77
United Kingdom	
Hungary	64
Guatemala	60
Philippines	57
Thailand	50
Italy (Sicily)	43
Severe iodine deficiency	
Marocco	27
Ethiopia	5-16
	64
Congo	15
	13

* Compiled from Semba-Delange 2001⁴¹ and Dorea 2002,⁴² where detailed data and references are to be found.

During lactation, considering that the iodine content of breastmilk in conditions of iodine sufficiency is in the range of 150-180 $\mu\text{g/L}$ ^{41,42} (Table 2) and that the milk production is from 0.5 to 1.1 liter per day up to the age of 6 months, the daily loss of iodine in human milk is estimated at some 75 to 200

$\mu\text{g/day}$. Consequently, the iodine requirement during lactation is estimated at 225 to 350 $\mu\text{g/day}$. The slight difference, if any, as compared to the figure of 290 $\mu\text{g/day}$ recommended by IOM¹² results from more recent data on the iodine content of breast milk.^{41,42}

Level of Urinary Iodine Indicating Optimal Iodine Nutrition During Pregnancy and Lactation

Considering that most (above 90%) of the iodine absorbed in the body eventually appears in the urine, urinary iodine excretion is a good marker of a very recent dietary iodine intake.¹¹ Therefore, a median urinary iodine in the general population varying from 100 to 199 $\mu\text{g/L}$ is considered as an indicator of an adequate iodine intake and an optimal status of iodine nutrition.¹¹ As the iodine requirement is increased during pregnancy, the median urinary iodine during pregnancy indicating optimal iodine nutrition needs to be higher than 100 $\mu\text{g/L}$. Table 1 compares the data available in the literature on urinary iodine in pregnant women and in the general population. In this Table, the countries are arbitrarily listed on the basis of roughly decreasing iodine intake of the general population, starting with Chile³¹ which is exposed to iodine excess based on the WHO/UNICEF/ICCIDD criteria,¹¹ down to countries where different degrees of mild to moderate iodine deficiency have been documented. As indicated earlier, there is a striking similarity between the urinary iodine in pregnant women and in the global population except in the reports published by Smyth et al.^{32,33} in which the values during pregnancy are systematically markedly higher than in non-pregnant controls. Therefore, it appears difficult to derive a reference value for urinary iodine during pregnancy and lactation from the data collected in countries with no iodine deficiency as this value varies from 800 $\mu\text{g/L}$ in Chile³¹ to 138 $\mu\text{g/L}$ in Switzerland, where the median urinary iodine in the

general population is barely above the lower limit of normal.³⁵ In Iran, where iodine deficiency has been successfully eliminated,⁴³ the median urinary iodine in pregnant women in four different cities varies from 186 to 403 $\mu\text{g/L}$ and is almost entirely similar to the values found in the general population in the same cities.⁴⁴ The values during pregnancy are of the same order of magnitude as the 250-300 $\mu\text{g/day}$ recommended as intake based on metabolic studies. And yet, in spite of these relatively elevated values, Azizi et al.⁴⁴ underline that with such medians, some 8% of the values are still below the critical threshold of 100 $\mu\text{g/L}$ for non-pregnant adults. They suggest that the recommended dietary intake of iodine during pregnancy should be still higher. It has to be recognized however, that this figure of 8% corresponds almost exactly to the percentage of values (7.2%) below the cut-off point of 50 $\mu\text{g/L}$ indicating at least moderate iodine deficiency in a general population when the median is between 100 and 200 $\mu\text{g/L}$.⁴⁵ This percentage is considered as acceptable⁴⁵ considering the well documented day to day variability of urinary iodine, including during pregnancy.⁴⁶⁻⁴⁹

From these different considerations, it can be concluded that the recommended median value for urinary iodine during pregnancy and lactation has to be based on theoretical grounds. If, as in non-pregnant adults, the recommended median (100 to 200 $\mu\text{g/L}$) corresponds to the recommended intake (150 $\mu\text{g/day}$), the median urinary iodine during pregnancy and lactation should be in the range 225-350 $\mu\text{g/L}$. If, on the contrary, this recommended median was based on a recommended intake of 225-350 $\mu\text{g/day}$ and a mean daily urinary volume of 1.5 L/day, it should be in the range of 150-230 $\mu\text{g/L}$, i.e. only slightly higher than the value recommended for non-pregnant adults.

It has to be recognized that thyroid function and volume remained perfectly normal during pregnancy in Iran⁴⁴ as well as in Chile³¹ for values still twice higher, which strongly suggests that these values are not excessive

and potential sources of side effects.^{50,51} On the contrary, in all countries submitted to some degree of iodine deficiency where the point has been investigated, thyroid function is critically impaired during pregnancy and in the neonate even when it remains normal in the general population.⁵²⁻⁵⁶ The anomalies include progressive decrease in free T_4 and increase in serum Tg and thyroid volume. The alterations are usually still more marked in the neonates than in the mothers.⁵² They are at least partly corrected by iodine supplementation during pregnancy and lactation.^{57,58} In summary, it appears that the recommended dietary intake of iodine during pregnancy (250-300 $\mu\text{g/L}$) and lactation (225-350 $\mu\text{g/L}$) should be higher than what has been proposed earlier, especially by WHO/UNICEF/ICCIDD,¹¹ and that a median urinary iodine indicating optimal iodine nutrition during pregnancy and lactation could be in the range 150-230 $\mu\text{g/L}$.

Requirement of iodine in Neonates

As underlined by the IOM,¹² no functional criteria of iodine status have been demonstrated that reflect response to dietary intake in infants. Consequently, the recommended intake of iodine in neonates reflects the observed mean iodine intake of young infants exclusively fed human milk in iodine replete areas. Up to the late sixties, the iodine content of breast milk in such areas was usually around 50 $\mu\text{g/L}$.^{41,42,59} Considering a daily intake of breast milk of some 0.6 to 1 liter in the neonate and young infant, the assumption was that an infant may get 30 to 50 $\mu\text{g/day}$ iodine in milk from an adequately fed mother.⁶⁰ However, it is well established that the iodine content of breastmilk is critically influenced by the dietary intake of the pregnant and lactating mother and of the general population and that much higher figures have been recorded more recently.^{41,42} Thus, again on theoretical grounds, the requirement of iodine in neonates was evaluated from metabolic studies by determining the value which

Table 3. Median or mean (in bold) urinary iodine (UI) concentrations ($\mu\text{g/L}$) in neonates in iodine sufficiency and iodine deficiency

Countries and location	n	Gestational age	Urinary iodine ($\mu\text{g/L}$) [*]	Range	Reference
Japan	118	FT Breastfed	736		Harada et al. 1994 ⁶⁵
Hokkaido	182	FT Bottlefed	521		
United States					
Boston	?	PT \leq 36 weeks	148	16-510	Brown et al. 1997 ⁸⁹
Torrance	50	FT	921		Delange et al. 1984 ⁹⁰
Canada					
Toronto	81	FT	148		Delange et al. 1986 ⁷²
The Netherlands					
Rotterdam	64	FT	162		Delange et al. 1986 ⁷²
Amsterdam	36	FT	150		Bakker et al. 1999 ⁹¹
Sweden					
Stockholm	39	FT	112		Delange et al. 1986 ⁷²
Stockholm	61	FT	96		Heidemann et al. 1984 ⁶⁵
Mild to moderate iodine deficiency					
Germany					
Nine towns 1983	461	FT	12-29		Heidemann et al. 1984 ⁶⁵
Berlin West 1985	87	FT	28		Delange et al. 1986 ⁷²
Kiel 1992	50	FT	33		Grebe et al 1993 ⁹²
Frankfurt 1992	21	FT	37		Bohles et al. 1993 ⁹³
Berlin West 1994	177	FT	31		Grüters et al. 1995 ⁹⁴
Berlin East 1994	213	FT	44		Grüters et al. 1995 ⁹⁴
Göttingen 2000	22	FT	50		Roth et al. 2001 ⁹⁵
Heidelberg 1999	32	FT	95		Klett et al. 1999 ⁹⁶
Belgium					
Brussels 1983	103	PT+FT	35	10-150	Delange et al. 1984 ⁹⁰
Brussels 1985	196	FT	48		Delange et al. 1986 ⁷²
Brussels 2000	90	FT	86		Ciardelli et al. 2001 ⁹⁷
Italy					
Rome 1985	114	FT	47		Delange et al. 1986 ⁷²
Catania 1985	14	FT	71		Delange et al. 1986 ⁷²
?towns 1995	195	FT	56	10-950	Rapa et al. 1996 ⁹⁸
Milano 1995	18	PT 30 weeks	123		Parravicini et al. 1996 ⁹⁹
Torino	9	FT	67	10-162	Bono et al 1998 ¹⁰⁰
France					
Lille 1985	82	FT	58		Delange et al. 1986 ⁷²
Toulouse 1985	37	FT	29		Delange et al. 1986 ⁷²
Ireland					
Belfast 1993	?	FT	100		Barakat et al. 1994 ¹⁰¹
Israel					
Tel Aviv 1996	55	PT 30-31 wks		55-100	Linder et al. 1997 ¹⁰²
Czech Republic					
Prague 1998	50	FT	79		Hnikova et al. 1999 ⁷⁰
Prisbram 1998	50	PT	78		
Hungary					
Budapest 2002	55	FT	35		Peter et al. 2003 ¹⁰³
Gyor 2002	65	FT	57		
Miskole 2002	54	FT	59		
Nyiregyhaza	35	FT	75		
Severe iodine deficiency					
Göttingen 1985	81	FT	15		Delange et al. 1986 ⁷²
Heidelberg 1985	39	FT	13		Delange et al. 1986 ⁷²
Freiburg 1985	39	FT	11		Delange et al. 1986 ⁷²

n: number; FT: Full-term, PT: Pre-term

* Values are medians or means (bold).

resulted in a situation of positive iodine balance, which is required in order to insure a progressively increased intrathyroidal iodine pool in the growing young infant. Such iodine balance studies were conducted in healthy preterm and in fullterm infants aged approximately one month in Belgium, a country with mild iodine deficiency.⁶¹ These studies, reported extensively elsewhere,⁶⁰ indicate that the iodine intake required in order to achieve a positive iodine balance is at least 15 µg/kg/day in fullterms and 30 µg/kg/day in preterms. This corresponds approximately to 90 µg/day and is consequently twice higher than the 1989 US recommendations of 40-50 µg/day⁶² but is still a bit lower than the present recommendation of 110 µg/day by the IOM.¹²

Level of Urinary Iodine Indicating Optimal Iodine Nutrition in Neonates

Table 3 summarizes the data from the literature on the median urinary iodine in neonates in countries or areas with iodine sufficiency and with different degrees of iodine deficiency. There is a large variability in the results even in iodine sufficient countries, where they vary from 736 µg/L in Hokkaido, Japan,⁶³ which is submitted to an extremely high iodine intake⁶⁴ to 96 µg/L in Stockholm.⁶⁵

Therefore, again, the data from the literature do not help substantially in identifying the optimal urinary iodine level and this level has also to be defined on the basis of theoretical considerations. Based on an iodine requirement of 90 µg/day and a volume of urines in neonates of some 0.4 to 0.5 liter/day,⁶⁶ the median urinary iodine indicating optimal iodine nutrition in neonates can be evaluated at some 180 to 225 µg/L when ignoring the fact that the iodine balance of the neonate should also be positive in order to constitute the iodine stores of the thyroid. This level, which is higher than the one recommended for schoolchildren and adults, is indeed observed when healthy young infants

are supplemented with a daily physiological dose of 90 µg/day.⁶⁷ It is also the value reported in some parts of the United States supposed to be iodine sufficient.^{68,69} On the other hand, studies reported in the literature in which urinary iodine has been determined simultaneously in mothers at delivery and in neonates during the first days of life^{39,70,71} indicate that these levels are almost similar in mothers and neonates. Therefore, based on the considerations on optimal urinary iodine in pregnant mothers, it can be extrapolated that the level in neonates should be around 150 to 230 µg/L, which is almost similar to the figure derived from the iodine requirements of the neonates.

The data reported from neonates in conditions of mild, moderate and severe iodine deficiency are indeed much lower than normal, down to less than 20 µg/L in Germany⁷² before the partly successful implementation of a program of voluntary salt iodization.⁷³ It is particularly interesting to observe that this level progressively increased with time in Germany and in Belgium for example following the implementation of programs of iodine supplementation^{73,74} and silent iodine prophylaxis, respectively.⁷⁵

In summary, the recommended dietary intake of iodine in neonates is 90 µg/day and the median urinary iodine to be expected when this requirement is met is 180 to 225 µg/L, a value almost similar to the one recommended for pregnant women.

Conclusion

Pregnant and lactating women and neonates are the main targets to the effects of iodine deficiency because of the impact of maternal, fetal and neonatal hypothyroxinemia on brain development of the progeny.⁵⁻¹⁰ Therefore, any program of salt iodization in a population should pay special attention to these particular groups. And yet, no firm recommendations are presently available on the level of urinary iodine indicating optimal iodine nutrition in these groups. This paper constitutes an attempt

to propose such normative values. It appears that an extensive review of the literature based in particular on the evaluation of urinary iodine in these groups in iodine replete populations does not offer clear answers to the questions because of the variability of individual results even in iodine sufficient countries. One first conclusion of this paper is thus that more accurate data should be collected in iodine sufficient countries, comparing systematically and at the same time the urinary iodine in the general population, in non-pregnant adults, schoolchildren, pregnant and lactating women and in neonates.

However, based on the data from the literature and on metabolic considerations, it is proposed that the recommended dietary intake of iodine is 250-300 µg/day for pregnant women, 225-350 µg/day for lactating women and 90 µg/day for neonates and young infants. It is proposed that the median level of urinary iodine indicating optimal iodine nutrition during pregnancy and lactation is in the range 150-230 µg/L. Recommendations for neonates are still more difficult not only because of the lack of accurate data but also because the neonate is not in a steady state regarding iodine metabolism and that urinary iodine probably represents a relatively imprecise estimation of the iodine intake. How-

ever, based on the data from the literature and on theoretical considerations, it can be concluded that the median urinary iodine indicating optimal iodine nutrition in the neonate should be in the same range of 180-225 µg/L, almost similar to the value recommended for their mothers.

It has to be emphasized again that these levels are higher than the ones recommended for the general population and are supposed to be potentially responsible for side effects in adolescents and non-pregnant adults.¹¹ Therefore, special attention should be focused on iodine supplementation and monitoring urinary iodine during pregnancy and possibly during the neonatal period in addition to programs of Universal Salt Iodization in countries with iodine deficiency.⁵⁸ This recommendation is particularly relevant considering that pregnant and lactating women and neonates have usually a limited access to salt in general and, consequently, to iodized salt, and that even in the United States, where the status of iodine nutrition is adequate in the general population (median urinary iodine of 145 µg/L), 6.7% of the pregnant women are still affected by moderate to severe iodine deficiency (urinary iodine below 50 µg/L).⁷⁶

References

1. Beckers C, Reinwein D. The Thyroid and Pregnancy. Stuttgart: Schattauer publishers; 1991: p.1-203.
2. Stanbury JB, Delange F, Dunn JT, Pandav CS. Iodine in pregnancy. New Dehli: Oxford University Press; 1998: p.1-297.
3. Glinioer D. The regulation of thyroid function in pregnancy: pathways of endocrine adaptation from physiology to pathology. *Endocr Rev.* 1997 Jun; 18(3):404-33.
4. Berghout A, Wiersinga W. Thyroid size and thyroid function during pregnancy. In: Stanbury JB, Delange F, Dunn JT, Pandav CS, editors. *Iodine in pregnancy.* Dehli: Oxford University Press; 1998: p. 35-54.
5. DeLong GR, Robbins J, Condliffe PG. Iodine and the brain. New York: Plenum Press; 1989: p.1-379.
6. Stanbury JB. The damaged brain of iodine deficiency. New York: Cognizant Communication Co; 1994: p.1-335.
7. Morreale de Escobar G, Obregon MJ, Escobar del Rey F. Is neuropsychological development related to maternal hypothyroidism or to maternal hypothyroxinemia? *J Clin Endocrinol Metab.* 2000 Nov;85(11):3975-87.
8. Delange F. Iodine deficiency as a cause of brain damage. *Postgrad Med J.* 2001 Apr;77(906):217-20.
9. Lavado-Autric R, Auso E, Garcia-Velasco JV, Arufe Mdel C, Escobar del Rey F, Berbel P, et al.

- Early maternal hypothyroxinemia alters histogenesis and cerebral cortex cytoarchitecture of the progeny. *J Clin Invest.* 2003 Apr;111(7):1073-82.
10. Zoeller RT. Transplacental thyroxine and fetal brain development. *J Clin Invest.* 2003 Apr;111(7):954-7.
 11. WHO, UNICEF, and ICCIDD. Assessment of the Iodine Deficiency Disorders and monitoring their elimination. Geneva: WHO publication. WHO/NHD/01.1; 2001: p. 1-107.
 12. Institute of Medicine, Academy of Sciences, USA. Dietary reference intakes for Vitamin A, Vitamin K, Arsenic, Boron, Chromium, Copper, Iodine, Iron, Manganese, Molybdenum, Nickel, Silicon, Vanadium and Zinc. Washington DC: National Academy Press; 2001: p. 1-773.
 13. Dworkin HJ, Jacquez JA, Beierwaltes WH. Relationship of iodine ingestion to iodine excretion in pregnancy. *J Clin Endocrinol Metab.* 1966 Dec;26(12):1329-42.
 14. Thomson CD. Dietary recommendations of iodine around the world. *IDD Newsletter.* 2002;18(3):38-42.
 15. Ladipo OA. Nutrition in pregnancy: mineral and vitamin supplements. *Am J Clin Nutr.* 2000 Jul; 72(1 Suppl):S280-90.
 16. Reinwein D, Jaspers C, Kirbas C, Zorlu A. Thyroxine substitution during pregnancy. In: C Beckers, D Reinwein, editors. *The Thyroid and Pregnancy.* Stuttgart: Schattauer publishers; 1991: p. 115-24.
 17. Glinoe D. The thyroid in pregnancy: a European perspective. *Thyroid Today.* 1995; 18:1-11.
 18. Glinoe D. The systematic screening and management of hypothyroidism and hyperthyroidism during pregnancy. *Trends in Endocrinol Metab.* 1998;9: 403-11.
 19. Mahillon I, Peers W, Bourdoux P, Ermans AM, Delange F. Effect of vaginal douching with povidone-iodine during early pregnancy on the iodine supply to mother and fetus. *Biol Neonate.* 1989; 56(4):210-7.
 20. Etling N. Concentration of thyroglobulin, iodine contents of thyroglobulin and of iodo-aminoacids in human neonates thyroid glands. *Acta Paediatr Scand.* 1977; 66:97-102.
 21. Costa A, De Filippis V, Panizzo M, Giraudi G, Bertino E, Arisio R, et al. Development of thyroid function between VI-IX month of fetal life in humans. *J Endocrinol Invest.* 1986 Aug;9(4):273-80.
 22. Delange F, Bourdoux P, Laurence M, Peneva L, Walfish P, Willgerodt H, et al. editors. Neonatal thyroid function in iodine deficiency. In: *Iodine deficiency in Europe. A continuing concern.* New York: Plenum Press; 1993: p.199-210.
 23. Savin S, Cvejic D, Nedic O, Radosavljevic R. Thyroid hormone synthesis and storage in the thyroid gland of human neonates. *J Pediatr Endocrinol Metab.* 2003 Apr-May;16(4):521-8.
 24. Beckers C. Iodine economy in and around pregnancy. In: C Beckers, D Reinwein, editors. *The Thyroid and pregnancy.* Stuttgart: Schattauer publishers; 1991: p. 25-34.
 25. Fisher DA. Clinical review 19: Management of congenital hypothyroidism. *J Clin Endocrinol Metab.* 1991 Mar;72(3):523-9.
 26. Van Vliet G. Neonatal hypothyroidism: treatment and outcome. *Thyroid.* 1999 Jan;9(1):79-84.
 27. Aboul-khair SA, Crooks J, Turnbull AC, Hytten FE. The physiological changes in thyroid function during pregnancy. *Clin Sci.* 1964 Oct;27:195-207.
 28. Wayne EJ, Koutras DA, Alexander WD. *Clinical aspects of iodine metabolism.* Oxford: Blackwell publishing; 1964.
 29. Dafnis E, Sabatini S. The effect of pregnancy on renal function: physiology and pathophysiology. *Am J Med Sci.* 1992 Mar;303(3):184-205.
 30. Lazarus JH, Kokandi A. Thyroid disease in relation to pregnancy: a decade of change. *Clin Endocrinol (Oxf).* 2000 Sep;53(3):265-78.
 31. Liberman CS, Pino SC, Fang SL, Braverman LE, Emerson CH. Circulating iodide concentrations during and after pregnancy. *J Clin Endocrinol Metab.* 1998 Oct;83(10):3545-9.
 32. Smyth PP, Hetherington AM, Smith DF, Radcliff M, O'Herlihy C. Maternal iodine status and thyroid volume during pregnancy: correlation with neonatal iodine intake. *J Clin Endocrinol Metab.* 1997 Sep;82(9):2840-3.
 33. Smyth PP. Variation in iodine handling during normal pregnancy. *Thyroid.* 1999 Jul;9(7):637-42.
 34. Kung AW, Lao TT, Chau MT, Tam SC, Low LC. Goitrogenesis during pregnancy and neonatal hypothyroxinaemia in a borderline iodine sufficient area. *Clin Endocrinol (Oxf).* 2000 Dec;53(6):725-31.
 35. Hess SY, Zimmermann MB, Torresani T, Bürgi H, Hurrell RF. Monitoring the adequacy of salt iodization in Switzerland: a national study of school children and pregnant women. *Eur J Clin Nutr.* 2001 Mar;55(3):162-6.
 36. Brander L, Als C, Buess H, Haldimann F, Harder M, Hanggi W, et al. Urinary iodine concentration during pregnancy in an area of unstable dietary iodine intake in Switzerland. *J Endocrinol Invest.* 2003 May;26(5):389-96.
 37. Bürgi H. Iodine deficiency in Switzerland. In: F Delange, A Robertson, E McLoughney, G Gerasimov, editors. *Elimination of Iodine Deficiency Disorders (IDD) in Central and Eastern Europe, the Commonwealth of Independent States, and the Baltic States.* Geneva: WHO publication. WHO/EURO/NUT/98.1; 1998: p. 15-20.
 38. Glinoe D, de Nayer P, Bourdoux P, Lemone M, Robyn C, van Steirteghem A, et al. Regulation of

- maternal thyroid during pregnancy. *J Clin Endocrinol Metab.* 1990 Aug;71(2):276-87.
39. Vermiglio F, Lo Presti VP, Finocchiaro MD, Battiato S, Grasso L, Ardita FV, et al. Enhanced iodine concentrating capacity by the mammary gland in iodine deficient lactating women of an endemic goiter region in Sicily. *J Endocrinol Invest.* 1992 Feb;15(2):137-42.
 40. Caron P, Hoff M, Bazzi S, Dufor A, Faure G, Ghandour I, et al. Urinary iodine excretion during normal pregnancy in healthy women living in the southwest of France: correlation with maternal thyroid parameters. *Thyroid.* 1997 Oct;7(5):749-54.
 41. Semba RD, Delange F. Iodine in human milk: perspectives for infant health. *Nutr Rev.* 2001 Aug; 59(8 Pt 1):269-78.
 42. Dorea JG. Iodine nutrition and breast-feeding. *J Trace Elem Med Biol.* 2002;16(4):207-20.
 43. Azizi F, Sheikholeslam R, Hedayati M, Mirmiran P, Malekafzali H, Kimiagar M, et al. Sustainable control of iodine deficiency in Iran: beneficial results of the implementation of the mandatory law on salt iodization. *J Endocrinol Invest.* 2002 May;25(5):409-13.
 44. Azizi F, Aminorroya A, Hedayati M, Rezvanian H, Amini M, Mirmiran P. Urinary iodine excretion in pregnant women residing in areas with adequate iodine intake. *Public Health Nutr.* 2003 Feb;6(1):95-8.
 45. Delange F, de Benoist B, Bürgi H. Median urinary iodine concentrations indicating adequate iodine intake at population level. *Bull WHO* 2002; 80:410-7.
 46. Rasmussen LB, Ovesen L, Christiansen E. Day-to-day and within-day variation in urinary iodine excretion. *Eur J Clin Nutr.* 1999 May;53(5):401-7.
 47. Als C, Helbling A, Peter K, Haldimann M, Zimmerli B, Gerber H. Urinary iodine concentration follows a circadian rhythm: a study with 3023 spot urine samples in adults and children. *J Clin Endocrinol Metab.* 2000 Apr;85(4):1367-9.
 48. Bürgi H, Bangerter B, Siebenhüner L. High day-to-day variability of urinary iodine excretion despite almost universal salt iodization in Switzerland. In: RM Geertman, editor. 8th World Salt Symposium. Amsterdam: Elsevier publishers; 2000: p.961-3.
 49. Thomson CD, Packer MA, Butler JA, Duffield AJ, O'Donoghue KL, Whanger PD. Urinary selenium and iodine during pregnancy and lactation. *J Trace Elem Med Biol.* 2001 Apr;14(4):210-7.
 50. Delange F, Lecomte P. Iodine supplementation: benefits outweigh risks. *Drug Saf.* 2000 Feb; 22(2):89-95.
 51. Braverman LE. Adequate iodine intake--the good far outweighs the bad. *Eur J Endocrinol.* 1998 Jul;139(1):14-5.
 52. Glinioer D, Delange F, Laboureur I, de Nayer P, Lejeune B, Kinthaert J, et al. Maternal and neonatal thyroid function at birth in an area of marginally low iodine intake. *J Clin Endocrinol Metab.* 1992 Sep;75(3):800-5.
 53. Berghout A, Endert E, Ross A, Hogerzeil HV, Smits NJ, Wiersinga WM. Thyroid function and thyroid size in normal pregnant women living in an iodine replete area. *Clin Endocrinol (Oxf).* 1994 Sep;41(3):375-9.
 54. Vermiglio F, Lo Presti VP, Scaffidi Argentina G, Finocchiaro MD, Gullo D, Squatrito S, et al. Maternal hypothyroxinaemia during the first half of gestation in an iodine deficient area with endemic cretinism and related disorders. *Clin Endocrinol (Oxf).* 1995 Apr;42(4):409-15.
 55. Eltom A, Eltom M, Elnagar B, Elbagir M, Gebre-Medhin M. Changes in iodine metabolism during late pregnancy and lactation: a longitudinal study among Sudanese women. *Eur J Clin Nutr.* 2000 May;54(5):429-33.
 56. Rotondi M, Amato G, Biondi B, Mazziotti G, Del Buono A, Rotonda Nicchio M, et al. Parity as a thyroid size-determining factor in areas with moderate iodine deficiency. *J Clin Endocrinol Metab.* 2000 Dec;85(12):4534-7.
 57. Chan S, Gittoes N, Franklyn J, Kilby M. Iodine intake in pregnancy. *Lancet.* 2001 Aug 18;358(9281):583-4.
 58. Zimmermann M, Delange F. Iodine supplementation of pregnant women in Europe: a review and recommendation. *Eur J Clin Nutr* 2004 Jul; 58(7):979-84.
 59. Delange F. Physiopathology of iodine nutrition. In: RK Chandra, editor. Trace Elements in nutrition of children. New York: Raven Press; 1985: p.291-9.
 60. Delange F. Requirements of iodine in humans. In: Delange F, Dunn JT, Glinioer D, editors. Iodine Deficiency in Europe. A continuing concern. New York: Plenum Press; 1993: p.5-16.
 61. Delange F. Iodine deficiency in Europe anno 2002. *Thyroid International.* 2002;5:1-19.
 62. National Research Council, Food and Nutrition Board. Recommended Dietary Allowances. Washington DC: National Academy Press; 1989: p. 213-7 and Table p. 285.
 63. Harada S, Ichihara N, Arai J, Honma H, Matsuura N, Fujieda K. Influence of iodine excess due to iodine-containing antiseptics on neonatal screening for congenital hypothyroidism in Hokkaido prefecture, Japan. *Screening* 1994;3:115-23.
 64. Suzuki H, Higuchi T, Sawa K, Ohtaki S, Horiuchi Y. Endemic coast goitre in Hokkaido, Japan. *Acta Endocrinol (Copenh).* 1965 Oct;50(2):161-76.
 65. Heidemann PH, Stubbe P, von Reuss K, Schurnbrand P, Larson A, von Petrykowski W. Iodine excretion and dietary iodine supply in newborn in-

- fants in iodine-deficient regions of West Germany Dtsch Med Wochenschr. 1984 May 18; 109(20):773-8. (German).
66. Behrman RE, VC Vaughan, and Nelson WE. Nelson Textbook of Pediatrics, 13th Ed. Philadelphia: W.B. Saunders publishing; 1987.
 67. Delange F, Wolff P, Gnat D, Dramaix M, Pilchen M, Vertongen F. Iodine deficiency during infancy and early childhood in Belgium: does it pose a risk to brain development? Eur J Pediatr. 2001 Apr; 160(4):251-4.
 68. Bryant WP, Zimmerman D. Iodine-induced hyperthyroidism in a newborn. Pediatrics. 1995 Mar; 95(3):434-6.
 69. Gordon CM, Rowitch DH, Mitchell ML, Kohane IS. Topical iodine and neonatal hypothyroidism. Arch Pediatr Adolesc Med. 1995 Dec; 149(12):1336-9.
 70. Hnikova O, Hromadkova M, Wiererova O, Bilek R. Follow-up study of iodine status in neonates and their mothers in 2 regions of the Czech Republic after a 3-year intervention Cas Lek Cesk. 1999 Apr 26;138(9):272-5. (Czech).
 71. Tajtakova M, Capova J, Bires J, Sebokova E, Petrovicova J. Thyroid volume, urinary and milk iodine in mothers after delivery and their newborns in iodine-replete country. Endocr Regul. 1999 Mar;33(1):9-15.
 72. Delange F, Heidemann P, Bourdoux P, Larsson A, Vigneri R, Klett M, et al. Regional variations of iodine nutrition and thyroid function during the neonatal period in Europe. Biol Neonate. 1986;49(6):322-30.
 73. Gartner R. IDD status in Germany. J Endocrinol Invest. 2003;26 (Suppl. to n° 9):2223.
 74. Meng W, Schindler A. Iodine supply in Germany. In: Delange F, Robertson A, McLoughney E, Gerasimov G, editors. Elimination of Iodine Deficiency Disorders (IDD) in Central and Eastern Europe, the Commonwealth of the Independent States, and the Baltic States. Geneva: WHO publication; WHO/EURO/NUT/98.1. 1998: p. 21-30.
 75. Delange F, Van Onderbergen A, Shabana W, Vandemeulebroucke E, Vertongen F, Gnat D, et al. Silent iodine prophylaxis in Western Europe only partly corrects iodine deficiency; the case of Belgium. Eur J Endocrinol. 2000 Aug;143(2):189-96.
 76. Hollowell JG, Staehling NW, Hannon WH, Flanders DW, Gunter EW, Maberly GF, et al. Iodine nutrition in the United States. Trends and public health implications: iodine excretion data from National Health and Nutrition Examination Surveys I and III (1971-1974 and 1988-1994) J Clin Endocrinol Metab. 1998 Oct;83(10):3401-8.
 77. Elnagar B, Eltom A, Wide L, Gebre-Medhin M, Karlsson FA. Iodine status, thyroid function and pregnancy: study of Swedish and Sudanese women. Eur J Clin Nutr. 1998 May;52(5):351-5.
 78. Soldin OP, Soldin SJ, Pezzullo JC. Urinary iodine percentile ranges in the United States. Clin Chim Acta. 2003 Feb;328(1-2):185-90.
 79. Barnett CA, Visser TJ, Williams F, Toor HV, Duran S, Presas MJ, et al. Inadequate iodine intake of 40% of pregnant women from a region of Scotland. J Endocrinol Invest. 2002. Suppl. to N° 7, Abstract P110:90.
 80. Kung AW, Lao TT, Low LC, Pang RW, Robinson JD. Iodine insufficiency and neonatal hyperthyrotopinaemia in Hong Kong. Clin Endocrinol (Oxf). 1997 Mar;46(3):315-9.
 81. Vermiglio F, Lo Presti VP, Castagna MG, Violi MA, Moleti M, Finocchiaro MD, et al. Increased risk of maternal thyroid failure with pregnancy progression in an iodine deficient area with major iodine deficiency disorders. Thyroid. 1999 Jan;9(1):19-24.
 82. Mocan MZ, Erem C, Telatar M, Mocan H. Urinary iodine levels in pregnant women with and without goiter in the Eastern Black Sea of Turkey. Trace Elements and Electrolytes. 1995;12:195-7.
 83. Pedersen KM, Laurberg P, Iversen E, Knudsen PR, Gregersen HE, Rasmussen OS, et al. Amelioration of some pregnancy-associated variations in thyroid function by iodine supplementation. J Clin Endocrinol Metab. 1993 Oct;77(4):1078-83.
 84. Nohr SB, Laurberg P, Borlum KG, Pedersen KM, Johannesen PL, Damm P, et al. Iodine deficiency in pregnancy in Denmark. Regional variations and frequency of individual iodine supplementation. Acta Obstet Gynecol Scand. 1993 Jul;72(5):350-3.
 85. Antonangeli L, Maccherini D, Cavaliere R, Di Giulio C, Reinhardt B, Pinchera A, et al. Comparison of two different doses of iodide in the prevention of gestational goiter in marginal iodine deficiency: a longitudinal study. Eur J Endocrinol. 2002 Jul;147(1):29-34.
 86. Romano R, Jannini EA, Pepe M, Grimaldi A, Olivieri M, Spennati P, et al. The effects of iodoprophylaxis on thyroid size during pregnancy. Am J Obstet Gynecol. 1991 Feb;164(2):482-5.
 87. Liesenkotter KP, Gopel W, Bogner U, Stach B, Gruters A. Earliest prevention of endemic goiter by iodine supplementation during pregnancy. Eur J Endocrinol. 1996 Apr;134(4):443-8.
 88. Mezosi E, Molnar I, Jakab A, Balogh E, Karanyi Z, Pakozdy Z, et al. Prevalence of iodine deficiency and goitre during pregnancy in east Hungary. Eur J Endocrinol. 2000 Oct;143(4):479-83.
 89. Brown RS, Bloomfield S, Bednarek FJ, Mitchell ML, Braverman LE. Routine skin cleansing with povidone-iodine is not a common cause of transient neonatal hypothyroidism in North America: a prospective controlled study. Thyroid. 1997 Jun;7(3):395-400.
 90. Delange F, Dalhem A, Bourdoux P, Lagasse R, Glinoeer D, Fisher DA, et al. Increased risk of pri-

- mary hypothyroidism in preterm infants. *J Pediatr*. 1984 Sep;105(3):462-9.
91. Bakker B, Vulmsa T, de Randamie J, Achterhuis AM, Wiedijk B, Oosting H, et al. A negative iodine balance is found in healthy neonates compared with neonates with thyroid agenesis. *J Endocrinol*. 1999 Apr;161(1):115-20.
 92. Grebe SF, Rebeski F, Gent J, Müller KD. Iodine balance in neonates and their mothers. *Klin Lab*. 1993;39:143-6.
 93. Bohles H, Aschenbrenner M, Roth M, von Loewenich V, Ball F, Usadel KH. Development of thyroid gland volume during the first 3 months of life in breast-fed versus iodine-supplemented and iodine-free formula-fed infants. *Clin Investig*. 1993 Jan;71(1):13-20.
 94. Grüters A, Liesenkotter KP, Willgerodt H. Persistence of differences in iodine status in newborns after the reunification of Berlin. *N Engl J Med*. 1995 Nov 23;333(21):1429.
 95. Roth C, Meller J, Bobrzik S, Thal H, Becker W, Kulenkampff D, et al. The iodine supply of newborns. Comparison of iodine absorption and iodine excretion of mother and child. *Dtsch Med Wochenschr*. 2001 Mar 23;126(12):321-5. (German).
 96. Klett M, Ohlig M, Manz F, Troger J, Heinrich U. Effect of iodine supply on neonatal thyroid volume and TSH. *Acta Paediatr Suppl*. 1999 Dec;88(432):18-20.
 97. Ciardelli R, Haumont D, Gnat D, Vertongen F, Delange F. The nutritional iodine supply of Belgian neonates is still insufficient. *Eur J Pediatr*. 2002 Oct;161(10):519-23.
 98. Rapa A, Chiorboli E, Corbetta C, Sacco F, Bona G. Study, A.U. Urinary iodine excretion (UIE) screening in newborns exposed to iodine-containing antiseptics. *Horm Res*. 1996;46:74.
 99. Parravicini E, Fontana C, Paterlini GL, Tagliabue P, Rovelli F, Leung K, Stark RI. Iodine, thyroid function, and very low birth weight infants. *Pediatrics*. 1996 Oct;98(4 Pt 1):730-4.
 100. Bona G, Chiorboli E, Rapa A, Weber G, Vigone MC, Chiumello G. Measurement of urinary iodine excretion to reveal iodine excess in neonatal transient hypothyroidism. *J Pediatr Endocrinol Metab*. 1998 Nov-Dec;11(6):739-43.
 101. Barakat M, Carson D, Hetherington AM, Smyth P, Leslie H. Hypothyroidism secondary to topical iodine treatment in infants with spina bifida. *Acta Paediatr*. 1994 Jul;83(7):741-3.
 102. Linder N, Davidovitch N, Reichman B, Kuint J, Lubin D, Meyerovitch J, et al. Topical iodine-containing antiseptics and subclinical hypothyroidism in preterm infants. *J Pediatr*. 1997 Sep;131(3):434-9.
 103. Peter F, Muzsnai A, Bourdoux P. Changes of urinary iodine excretion of newborns over a period of twenty years. *J Endocrinol Invest*. 2003;26(2 Suppl):S39-42.